

RECEIVED: March 28, 2018

REVISED: May 7, 2018

ACCEPTED: June 20, 2018

PUBLISHED: June 28, 2018

Yang-Baxter deformations beyond coset spaces (a slick way to do TsT)

I. Bakhmatov,^{a,b} E. Ó Colgáin,^{a,c} M.M. Sheikh-Jabbari^d and H. Yavartanoo^e

^aAsia Pacific Center for Theoretical Physics,
Postech, Pohang 37673, Korea

^bInstitute of Physics, Kazan Federal University,
Kremlevskaya 16a, 420111, Kazan, Russia

^cPhysics Department, Postech,
Pohang 37673, Korea

^dSchool of Physics, Institute for Research in Fundamental Sciences (IPM),
P.O. Box 19395-5531, Tehran, Iran

^eState Key Laboratory of Theoretical Physics, Institute of Theoretical Physics,
Chinese Academy of Sciences, Beijing 100190, China

E-mail: ilya.bakhmatov@apctp.org, ocolgain.eoin@apctp.org,
jabbari@theory.ipm.ac.ir, yavar@itp.ac.cn

ABSTRACT: Yang-Baxter string sigma-models provide a systematic way to deform coset geometries, such as $AdS_p \times S^p$, while retaining the σ -model integrability. It has been shown that the Yang-Baxter deformation in target space is simply an open-closed string map that can be defined for any geometry, not just coset spaces. Given a geometry with an isometry group and a bivector that is assumed to be a linear combination of antisymmetric products of Killing vectors, we show the equations of motion of (generalized) supergravity reduce to the Classical Yang-Baxter Equation associated with the isometry group, proving the statement made in [1]. These results bring us closer to the proof of the “YB solution generating technique” for (generalized) supergravity advertised in [1] and in particular provide an economical way to perform TsT transformations.

KEYWORDS: Classical Theories of Gravity, Integrable Field Theories, Supergravity Models

ARXIV EPRINT: [1803.07498](https://arxiv.org/abs/1803.07498)